

PUBLIKATIONEN

»The healing of free gingival autografts for socket seal surgery – a case report«,
Thalmair T., Hinze M., Bolz W., Wachtel H. Eur J Esthet Dent 2010; 5(4): 358-368.

ABSTRACT

Tooth extraction will be followed by marked alterations to the tissue volume, in particular in the anterior zone, which may jeopardize the esthetic outcome of any dental treatment involving tooth extraction. If, however, ridge collapse can be prevented or minimized after tooth extraction, more predictable outcomes with superior esthetics can be accomplished along with fewer surgical procedures. Therefore, it was proposed that stabilizing the soft tissue architecture with a free gingival graft could minimize the soft tissue shrinkage. The following case report describes some surgical modifications and refinements in order to enhance the predictability of the socket-seal surgery.

Für weitere Informationen zu dieser Publikation kontaktieren Sie
uns bitte unter den u. g. Kontaktdataen.
Die unautorisierte Verwendung dieses Textes ist nicht gestattet;
alle Rechte vorbehalten.

Dr. Marc Hinze
Spezialist für Implantologie & Parodontologie

Bahnhofstraße 17 82166 Gräfelfing Tel.: +49 (0)89 851076 Fax: +49 (0)89 8543559
info@marc-hinze.de www.marc-hinze.de